



The Road to Functional Bioanalysis: Development and Validation of a Cell-Based Assay for Neutralizing Anti-Drug Antibody Analysis

**Christelle Pythoud, PhD
Celerion Switzerland AG**

**2nd EBF YSS, Barcelona Spain
Nov 17, 2015**

Outline


- Introduction
 - Immunogenicity and functional bioanalysis
 - Basics of cell-based assays
 - The use of cell-based assays
- Development of a CBA for NAb testing: challenges and outcomes
 - Choose the relevant assay format and endpoint
 - Choose the relevant cell line and readout
 - Cell bank establishment
 - G-CSF activity and neutralization assay
 - Matrix interference assay – Specificity assay
- Validation Steps of a Cell-based NAb Assay
- Conclusion

Immunogenicity and Functional Bioanalysis

- Immune response to therapeutic proteins ► patient safety and drug efficacy issues
- Major safety concern: **neutralizing antibodies (NAbs)**, able to neutralize the biological activity of bound Ag
 - ► loss of product efficacy by binding to the product active site (highly critical if the product is a lifesaving drug)
 - ► cross-reactivity to and inhibition of the endogenous counterpart of the therapeutic protein (highly critical if the endogenous protein is nonredundant)
- The demand for drug immunogenicity assessment and characterization is growing (2014 FDA Guidance on “Immunogenicity Assessment for Therapeutic Protein Products”)
- Key safety data that impact critical decisions on the continuation of a drug development project
- **Cell-based assays (CBA)** (also called bioassay) are recognized by regulatory authorities as the gold standard to measure drug propensity to generate NAbs

Basics of Cell-based Assays

The 3 stages of the cell-signaling process



- The types of response are highly diverse, all measurable
- The 3 different steps of cell-signaling are individually measurable
- **The cell is a test system that allows a vast number of assay possibilities measuring a biological activity**
 - ▶ **Functional Bioanalysis**

Use of Cell-based Assays

- Determination of biologics potency for lot release
- Screening drug candidate targeting specific biological process during drug discovery
- **Functional bioanalysis for neutralizing antibody testing**

| | CBA | Non-CBA |
|------|--|--|
| PROs | <ul style="list-style-type: none">■ Test for the functionality/biological activity■ Closer to physiological conditions■ Recommended by authorities for NABs assessment■ More complex: more assay format and readout possibilities | <ul style="list-style-type: none">■ Less variability■ Easier to develop■ Faster to develop |
| CONS | <ul style="list-style-type: none">■ Higher variability■ More complex: experienced assay designer required, time-consuming | <ul style="list-style-type: none">■ No functional characterization |



Development of a CBA for NAb Testing: Challenges and Outcomes

Development of a CBA for NAb Testing

1. Choose the relevant assay format and endpoint


- The assay format is derived from the drug molecular mechanism
- Understanding the biological activity of the drug is essential to selecting the most appropriated cell-based assay format and endpoint
- What are the critical biological pathways involved and can they be exploited to develop a bioassay?

▶ *Development of a CBA to detect NAb against a PEGylated form of human G-CSF*


Biological Activity and Molecular Mechanism of G-CSF

- **G-CSF = Granulocyte - Colony Stimulating Factors**
- In vivo, G-CSF stimulates the bone marrow to induce differentiation into and proliferation of neutrophil granulocytes (aka neutrophils)
- Neutrophils: most abundant white blood cells (70%), essential part of the innate immune system, one of the first cells to migrate towards the site of infection to digest bacteria, major component of pus
- In clinical setup, recombinant human G-CSF (Filgrastim) is used to stimulate neutrophils production in patients suffering from neutropenia (low neutrophil counts, congenital or chemotherapy-induced)

Biological Activity and Molecular Mechanism of G-CSF



► *Cell proliferation assay to detect NAbs against G-CSF*



► *Direct neutralization assay (inhibition of stimulation), measures the inhibition of the drug activity*


Development of a CBA for NAb Testing

2. Choose the relevant cell line and readout

- The assay format conditions the choice of the cell line and readout
 - ▶ *NFS-60 cell line with alamar blue test as readout for cell proliferation (measure of the cell metabolic activity, proportional to number of cells at a defined time-point, fluorescence readout)*

Confirmation of NFS-60 phenotype

- NFS-60 cells express G-CSFR and IL3R
- Proliferation in response to hG-CSF and mIL3



Cell Bank Establishment

Origin/Source

- Cell background/species
- History, genetic modifications
- Number of passages
- Cell culture conditions

Documentation

- Cell background/species
- History, genetic modifications
- Number of passages
- Cell culture conditions
- Biosafety

Characterization of CB

- Viability
- Growth curve



Biosafety

- BSL2 conditions
- Bacterial contamination free


Cell bank (CB)

- Tiered cell bank system
(Initial CB → Master CB → Working CB)
- (identical passage number at each level)
- Qualified storage container
(vapor phase of liquid N₂)

G-CSF Activity and Neutralization Assay



Cells + G-CSF
 ↓ 72h
 cell number measurement




Cells + G-CSF/NAb anti-G-CSF
 ↓ 72h
 cell number measurement


Matrix Interference Assay – Specificity Assay

- More than any other method, CBA performance can be affected by factors present in matrix
- Matrix interference assay is recommended in NAb testing
- Differentiation between drug-specific NAb and other non-specific inhibitory factors**

- *Alternative stimulus assay*
 - Induction of cell proliferation using another stimulus: *mIL3*
 - Anti G-CSF NAb are not able to inhibit *mIL3*-induced NFS-60 cell proliferation ► **assay specificity**



Cells + mIL3
 ↓ 72h
 cell number measurement



Cells + mIL3/NAb anti-G-CSF
 ↓ 72h
 cell number measurement

Validation Steps of a Cell-based NAb Assay

- No regulatory guidelines, but industry standard:
 - Gupta S. et al., Recommendations for the design, optimization, and qualification of cell-based assays used for the detection of neutralizing antibody responses elicited to biological therapeutics, *Journal of Immunological Methods* (321) 1-18, 2007
- Cut point calculation (10-20 drug naïve healthy individuals, statistical approach)
- Sensitivity
- Selectivity in individuals
- Specificity/matrix interference assay
- Free drug tolerance
- Precision
- Stability of PC antibodies
- Robustness (determination of acceptance criteria for cell passage number)
- Hemolysis

Conclusion

- Validation steps for cell-based NAb assay are similar to those for binding ADA assay
- Living cells as test system
- “Reagent” more difficult to control as compared to chemicals
- Large contributor to assay variability
- Requirement for a controlled process to ensure continuous availability of a consistent and reliable cell source ► cell banking
- Requirement for specificity assay performed in parallel to the neutralizing assay ► matrix interference assay
- Understanding the drug’s mode of action as well as some of the clinical features (study population) is critical for cell-based method development

Acknowledgments

Guy Lemaillet, PhD

Christoph Köhler

Petra Struwe, PhD

Christine Schiebl, PhD

***Come and see our posters
at Celerion Booth #B25***



Thank you for your attention